

COVID-19: DOSE PER POUND FOR PROPHYLAXIS PRE-EXPOSURE WITH IVERMECTIN FOR EXPOSED

DOSE: 0.091 MG. PER POUND OF WEIGHT. 2nd DOSE IN 1 WEEK (7 D). THEN REPEAT DOSE EVERY 2 WEEKS IN THOSE WHO ARE EXPOSED TO PLACES WITH A HIGH VIRAL LOAD (HOSPITALS, MARKETS, TERMINALS OR SIMILAR) AND THEIR “CONTACTS” THAT ARE MEN > 50 AND WOMEN > 75, REPEAT THE DOSE EVERY WEEK (7 D)

PRESENTATIONS AND EQUIVALENCES FOR EACH DOSE OF 0.091 mg. PER POUND OF WEIGHT or 0.2 mg. PER KILO

WEIGHT		DOSE X WEIGHT	TABLETS of 3 mg. (if they are 6 mg. give half)	BOTTLE at 1% (10 mg. per ml.)	ORAL PASTE at 1.87%, with blue plunger (2.27 mg. per every: Notch, Teeth or 1/16”)	
Kilos	Libras					
	X	X lb x 0.091	X lb between 33 = tab. to give	X lb per 0.0091 = ml.	X lb x 0.04 = N° notch and teeth, or N° 1/16”	
34.0	75	6.8 mg	2 tablets	0.7 ml	3: 2 notch and 1 teeth	3/16”
40.8	90	8.2 mg	3 tablets	0.9 ml.		
45.4	100	9.1 mg	3 tablets	0.9 ml.	4: 2 notch and 2 teeth	4/16”= 1/4“
49.9	110	10.0 mg	3 tablets	1.0 ml.		
56.7	125	11.4 mg	4 tablets	1.1 ml.	5: 3 notch, 2 teeth: 1/2 measure	5/16”
63.5	140	12.7 mg	4 tablets	1.3 ml.		
68.0	150	13.7 mg	5 tablets	1.4 ml.	6: 3 notch and 3 teeth	6/16”= 3/8”
74.8	165	15.0 mg	5 tablets	1.5 ml.		
79.4	175	15.9 mg	5 tablets	1.6 ml.	7: 4 notch and 3 teeth	7/16”
83.9	185	16.8 mg	6 tablets	1.7 ml.		
90.7	200	18.2 mg	6 tablets	1.8 ml.	8: 4 notch and 4 teeth	8/16”= 1/2“
95.3	210	19.1 mg	7 tablets	1.9 ml.		
102.1	225	20.5 mg	7 tablets	2.1 ml	9: 5 notch and 4 teeth	9/16”
106.6	235	21.4 mg	7 tablets	2,2 ml		
113.4	250	22.8 mg	8 tablets	2.3 ml	10: 5 notch, 5 teeth.: 1 measure	10/16”= 5/8”

With the Bottles it is recommended to use a 5 ml syringe to measure the doses more accurately. Take it after lunch or dinner. Do not take with fruit juice, lemonade or milk. In addition to Ivermectin you should take a Multivitamin that contains Zinc, Vitamin D, C and A, or give these individually. It is indicated to gargle and nasal washings post-exposure to places of high viral load.

Source: Gustavo Aguirre Chang; Aurora Trujillo F. COVID-19: Prophylaxis Pre-Exposure with Ivermectin in exposed people. July 21, 2020.